

RESEARCH

Open Access



The effect of a simulation-based training program in basic life support on the knowledge of Palestinian nurses: a quasi-experimental study in governmental hospitals

Yousef Fahajan¹, Osama J. Emad², Ahmed Hassan Albelbeisi^{3,4}, Ali Albelbeisi⁵, Yasmin Abu Shnena⁶, Ayda Khader¹ and Edris Kakemam^{7*}

Abstract

Background Basic Life Support (BLS) plays an important role in increasing the survival rate of hospitalized heart attack patients. There are no previous studies on the effect of BLS training among Palestinian nurses. This study aimed to evaluate the effect of simulation-based BLS training program on nurses' knowledge Palestinian nurses at governmental hospitals.

Methods A quasi-experimental, pre & post-test design was used. 700 nurses were recruited proportionally using a simple random sampling method among 2980 nurses from 13 public hospitals in the Gaza Strip. This study was conducted from June to August 2022. A practical BLS test consisting of 10 multiple-choice questions according to American Heart Association guidelines (2020) was collected and sociodemographic characteristics. SPSS software, version 24 was used for the statistical analysis. Descriptive statistics and weighted mean were used. T-Test and One-way analysis of variance (ANOVA) were applied to determine differences in means among groups.

Results Most of the participating nurses (55.7%) were male, while (44.3%) were female. The majority of nurses (84.4%) are under 40 years of age. The weighted mean scores in the pre-test ranged from 52.2 to 75.1% and the mean scores was (6.16 ± 1.97). After applying conventional BLS training, the weighted mean scores ranged from 85.6 to 97.3% and the mean scores was (9.19 ± 1.04). The study revealed that the nurses' knowledge increased after applying simulation-based training program. The mean of knowledge scores was statistically significant between the pre and post-test on the basis of the current work hospital (P-value < 0.001).

Conclusion This study affords significant evidence of the positive effects of the BLS training program in improving nurses' knowledge; we recommend advanced BLS training for all healthcare providers, doctors, and nurses working in hospitals and healthcare centers. Nursing managers can implement systematic strategies to enhance nurses' knowledge and practice in BLS to target low-scoring Governorates.

Keywords Simulation, BLS Knowledge, BLS Practice, Nurses

*Correspondence:

Edris Kakemam
edriskakemam@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

A heart attack is an unexpected cessation of the functional activity of the heart due to failure of the heart to pump blood successfully [1]. Heart attack as a life-threatening state needs urgent interventions to ensure the lives of individuals and avoid irreversible damage to vital organs. According to the American Heart Association (AHA), cardiovascular disease has been identified as the number one cause of death, accounting for more than 17.9 million deaths per year in 2015, a number that is anticipated to extend to more than 23.6 million by 2030 [2]. A heart attack a major health problem that is estimated to be responsible for 15–20% of all deaths [3]. Several studies showed that the cardiac arrest survival rate decreases by 5–10% for each minute within the frame of cardiopulmonary resuscitation, which is an integral component of (BLS) [4]. BLS serves as a critical determinant of patient survival from heart attack. Training is vital in the enhancement of Cardiopulmonary Resuscitation (CPR) effects. BLS includes recognition, support of ventilation, and circulation in the episode of heart attack. It has a mixture of abilities including mouth-to-mouth breathing to support ventilation and chest compressions to normalize blood circulation to the brain and vital organs, called (CPR). BLS may be a basic and very effective method that allows anyone to help assist maintain life, soon after heart attack [5].

Knowledge and practice of nurses regarding BLS strategies ensure that the patient survives long sufficient [6]. Nurses' knowledge of BLS and proper practice of the techniques and maneuvers enable a person to effectively resuscitate a patient and its importance in saving lives [7]. Simulation-based BLS training is a viable way of exchanging abilities with trainees in a cost-effective way. It provides an optimum way for employers to assess how well their trainees are putting abilities into practice, and the decisions they are making in front of simulated real-life circumstances [8]. Using simulation-based BLS training ultimately extends the quality of treatment and patient safety through a greater commitment by healthcare professionals to principles-based resuscitation approaches [9]. Also, the use of the simulation approach can increase the knowledge, skills, and performance of the nursing practice; they can reach high levels of critical thinking and obtain new professional skills without putting the patient's health at risk [10]. A previous systematic review study aimed to evaluate effect of simulation-based training on nurses' skills and knowledge, demonstrated that the simulation training appears to be an effective strategy for improving nurses' knowledge and skills [11]. In addition, a qualitative study conducted among midwifery students in Sweden showed that simulation created links between theory and practice, and provided a safe learning environment [12]. To the best of our knowledge, there

are no previous studies about BLS training in Palestine. Therefore, this study aimed to evaluate the effect of simulation-based BLS training on nurses' knowledge among nurses at governmental hospitals in the Gaza strip, Palestine.

Methods

Design and setting

A quasi-experimental design, pre & post-test conducted from June to August 2022 in 13 governmental hospitals located in the five Gaza Strip Governorates' Gaza Strip is a small portion of the Occupied Palestinian Territories and is divided into five small governorates (Rafah, Khan-Younis, Middle-zone, Gaza, and North-Gaza) with an inhabitant of more than two million [13, 14]. The included hospitals have various services and units such as pediatrics, surgery, eye care, psychiatry, and pediatric care.

Participants

The study targeted of 2980 licensed nurses employed in 13 government hospitals.

Inclusion criteria All nurses should meet the following criteria: (1) nurses have at least a diploma degree; (2) employed in the 13 governmental hospitals (3) work experience of at least 6 months, (3) and fixed nurses.

Exclusion criteria Newly appointed nursing staff, part-time nurses, and the nurses have administrative roles were excluded.

Sample size and sampling method

The sample size for this study was estimated at 786 based on Krejci and Morgan's table (margin of error=3%, confidence level=95%). In fact, when the total population under study is known, the simplest and easiest way to estimate the study's sample size is the Morgan Table [15]. Hence, 786 male and female nurses in the surveys were proportionately selected based on the number of nurses in each hospital. A simple random sampling method was used to select participants from each hospital.

Intervention

The conventional adult BLS training was carried out by BLS-qualified trainers in collaboration with the head of the in-service health education department at each hospital. PowerPoint, movies, and hands-on exercises on CPR mannequins were used. The training was done for two hours per week for one month for each group, Before the start of the training, the pre-test was conducted, and after the end of the training, the post-test was conducted, the participants continued to practice various BLS roles including heart compression, artificial respiration, the

recovery position, airway management, foreign body airway obstruction, Administer first aid to an unconscious patient, and use of the Automated External Defibrillator (AED). Each participant practiced BLS stages for 5 min and at least of 20 min in each group after practicing on mannequins under the supervision of the trainers. Each subject is practiced independently and taught as a critical skill. The training was mainly based on role-playing and simulation, with all groups implementing the same training methods.

Measures and data collection

Before the start of the simulation-based training program, a pre-test was distributed and collected by a qualified, trained nurse in each hospital, and every questionnaire with missing data was excluded. After completion of the simulation-based training program, a post-test was distributed and collected by a qualified, trained nurse at each hospital, and every questionnaire with missing data was excluded. The questionnaire included three variables related to the participant's age, gender, and current work hospital. In addition, the BLS practice test contains of 10 multiple-choice questions and adheres to the AHA (2020) guidelines [16, 17]. The questionnaire includes items regarding compression-to-ventilation ratio, correct steps to operate an AED, pulse checking and correct locations, BLS steps used for adults, vital characteristics of first-rate CPR, recommended BLS sequence for the 2020 ILCOR guidelines, and signs of

an obstruction of the airway. The overall score for this tool was from 0 to 10, and a higher score revealed more knowledge about BLS.

Data analysis

SPSS software, version 24 was used for the statistical analysis. The characteristics of the nurses were described by using descriptive statistics such as mean, standard deviation, and percentages. In addition, weighted mean was calculated, calculating weighted mean involves multiplying each data point by its weight and summing those products. Then sum the weights for all data points. Finally, divide the weight*value products by the sum of the weights [18]. T-Test and One-way analysis of variance (ANOVA) were applied to determine differences in means among groups. All tests were conducted at the 0.05 level of statistical significance.

Results

Characteristics of participants

Of the 786 questionnaires distributed, 700 questionnaires were analyzed (response rate: 89.1%). The results of this study revealed that, more than half of the nurses (55.7%) were male, while (44.3%) were female. The majority of nurses (84.4%) are under 40 years of age. Almost a third of the participants were from Al-Shifa medical complex, while the least number of participants were from the Psychiatric Hospital and Al- Naser Eye Hospital (Table 1).

Knowledge of the study participants before and after the training

Table 2 displays the weighted mean scores of nurses to the BLS practice test questions according to the AHA (2020) guidelines. The weighted mean score in the pre-test ranged from 52.2 to 75.1%, while in the post-test they ranged from 85.6 to 97.3%. The nurses' weighted mean score was higher on the post-test in the 10 questions included.

Knowledge Mean scores before and after training

Table 3 displays that the mean knowledge was statistically significant between the pre and post-test. Nurses' mean scores in the pre-test was (6.16 ± 1.97) and the mean score in the post-test was (9.19 ± 1.04) .

SD: Standard Deviation. Differences between means were tested by using the Paired t-test.

Knowledge mean scores before and after training based on gender

Table 4 displays that the mean scores of knowledge were non-statistically significant between the pre and post-test on the basis of gender. The mean scores for the pretest were (06.21 ± 1.96) and (06.18 ± 1.98) for males and females, respectively ($p=0.361$). The mean scores for the pretest

Table 1 Demographic characteristics of the nurses (n = 700)

Variable	Frequency	Percentage %
Gender		
Male	390	55.7
Female	310	44.3
Age(years)		
20-30years	273	39.0
31-40 years	318	45.4
41-50 years	81	11.6
51-60 years	28	4.0
Current work hospital		
Kamal Odwan	17	2.4
Al-Naser Eye Hospital	12	1.7
Al-Najar	35	5.0
Psychiatric	12	1.7
Beit-Hanon	24	3.4
Al-Shifa Complex	207	29.6
Naser Complex	115	16.4
Al-Emirate	31	4.4
Indonesia	54	7.7
European Gaza	93	13.3
Al-Dora	21	3.0
AL-Rantisi	62	8.9
Turkish	17	2.4

Table 2 Knowledge of the study participants before and after the training

Question	Pretest	Post-test
	Weighted mean (%)	Weighted mean (%)
What is the compression-to-ventilation ratio you should use when giving CPR to any individual?	52.2	93.1
When performing two-rescuer CPR, how often should you switch roles?	52.7	88.1
When operating an AED, what are the correct steps to follow?	61.4	85.6
When looking for a pulse on a child from one year to puberty, where should you check?	69.4	92.4
What are the BLS (Basic Life Support) steps used for adults?	52.3	93.6
What are the vital characteristics of first-rate CPR?	75.1	97.3
Which step is NOT a part of the five steps in the Adult Chain of Survival?	57.4	93.6
What is the recommended BLS sequence for the 2020 ILCOR guidelines?	66.6	94.4
What are the signs of an obstruction of the airway?	68.9	92.6
How many breaths should be given during a two-rescuer CPR on an adult with an advanced airway in place?	60.1	89.1
Overall Weighted mean score (%)	61.6	91.9

Table 3 Knowledge mean score before and after training

	Mean	SD	P-value
Pre-Test	06.16	1.97	<0.001
Post-Test	09.19	1.04	

Table 4 Knowledge mean scores before and after training based on gender

Gender (N)	Mean	SD	P-value	
Pre-Test	Male (390)	06.21	1.96	0.361
	Female (310)	06.08	1.98	
Post-Test	Male (390)	09.19	1.06	0.948
	Female (310)	09.20	1.01	

Differences between means were tested by using the Independent-samples t-test

Table 5 Knowledge mean scores before and after training based on age groups

Age group/Years (N)	Mean	SD	P-value	
Pre-Test	20–30(273)	06.06	02.11	0.725
	31–40 (318)	06.24	01.91	
	41–50 (81)	06.08	01.81	
	51–60 (28)	06.17	01.70	
Post-Test	20–30(273)	09.20	01.07	0.343
	31–40 (318)	09.13	01.04	
	41–50 (81)	09.34	00.98	
	51–60 (28)	09.35	00.78	

Differences between means were tested by using the One-Way ANOVA

were (09.19 ± 1.06 and 09.20 ± 1.01) for males and females, respectively ($p=0.948$).

Knowledge mean score before and after training based on age groups

Table 5 displays that the mean scores of knowledges were non-statistically significant between the pre and post-test on the basis of age group. The highest mean score in the pretest was registered in the age group 31–40 years old (06.24 ± 1.9) and the lowest mean score was registered in the age group 20–30 years old (06.06 ± 2.11). The

highest mean score in the post-test was registered in the age group 51–60 years old (09.35 ± 0.78) and the lowest mean score was registered in the age group 31–40 years old (09.13 ± 1.04).

Knowledge mean scores before and after training based on the current work hospital

Table 6 shows that the mean scores of knowledges were statistically significant between the pre and post-test on the basis of the current work hospital (P -value<0.001). In the pre-test, the highest mean score was registered in the Turkish Hospital located in Gaza governorate (08.35 , $SD \pm 1.04$). The lowest was in Kamal Odwan Hospital (04.64 , $SD \pm 2.20$) located in North-Gaza governorate. In the post-test, the highest mean score was registered in Al-Dora Hospital located in Gaza governorate (10.00 , $SD \pm 0.20$). The lowest was registered in Kamal Odwan Hospital (07.58 , $SD \pm 1.32$) located in North-Gaza governorate.

Discussion

A simulation training is a dynamic method used to correlate actual clinical conditions in a safe area that supports nurses to grow knowledge and psychomotor skills resuscitation [19]. To the best of our knowledge, there are no previous studies about BLS training on nurses' knowledge in Palestine. Therefore, the study aimed to evaluate the effect of simulation-based BLS training on nurses' knowledge among nurses at public hospitals in the Gaza strip, Palestine.

The results of this study showed that the weighted mean scores in the pre-test ranged from 52.2 to 75.1% and the mean score was (6.16 out of 10.0 ± 1.97). Although our participants had a mean score higher than the mean score in a study conducted among nurses in Turkey (5.66 ± 1.97) [20]. The current findings indicate the need for urgent intervention to enhance the knowledge of Palestinian nurses in BLS. It has been stated that

Table 6 Knowledge mean scores before and after training based on the current work hospital

Current work hospital	Pre-test			Post-test		
	Mean	SD	P-value	Mean	SD	P-value
Kamal Odwan (17)	04.64	2.20	< 0.001	07.58	1.32	< 0.001
Al-Shifa Complex (207)	06.20	2.0		09.14	1.09	
Naser Complex (115)	06.35	1.71		09.12	0.98	
Al-Elmirate (31)	06.25	1.73		09.51	0.72	
Indonesia (54)	06.24	2.17		09.11	0.94	
European Gaza (93)	05.94	2.12		09.65	0.56	
Al-Dora (21)	05.23	1.41		10.00	0.20	
AL-Rantisi (62)	05.45	1.87		09.25	0.95	
Turkish (17)	08.35	1.49		09.11	1.05	
Al- Naser Eye (12)	07.33	1.49		07.91	1.31	
Al-Najar (35)	06.57	1.37		09.31	0.75	
Psychiatric (12)	05.50	1.73		08.16	1.64	
Beit-Hanon (24)	06.58	2.06		09.29	1.04	

Differences between means were tested by using the One-Way ANOVA

BLS is a dynamic issue that needs continuous and simultaneous training. There is a relationship between BLS knowledge, nurse practice, and cardiopulmonary resuscitation (CPR) consequences [21]. After applying conventional BLS training for adults by qualified BLS instructors and participants practiced BLS steps in the Clinical Skills Center, the weighted mean scores in post-test they ranged from 85.6 to 97.3% and the mean score was (9.19). The study displayed that the nurses' knowledge enhanced after implementing simulation-based training. These outcomes are consistent with earlier studies that showed a significant effect of CPR training on knowledge and practice [22–25]. The results of a study conducted among undergraduate nursing students in Thailand showed that the training had a significant and immediate impact on knowledge, self-efficacy, and chest compression skill [22]. Another study compared the knowledge and skills of undergraduate and staff nurses showed that the knowledge and skills of both groups of nurses improved with training [23].

The current study findings showed that the mean scores of knowledges were non-statistically significant between the pre and post-test on the basis of gender and age group. Similar results were found in a study carried out among nurses in Kuwait hospitals [26]. The mean scores of knowledges were statistically significant between the pre and post-test on the basis of the current work hospital (P -value < 0.001). In the pre-test, the highest mean score was registered in the Turkish Hospital located in Gaza governorate. The lowest was in Kamal Odwan Hospital located in North-Gaza governorate. In the post-test, the highest mean score was registered in Al-Dora Hospital located in Gaza governorate. The lowest was registered in Kamal Odwan Hospital located in North-Gaza governorate. The current findings can be explained by the fact that the skills of health care workers'

in some governorates of the Gaza Strip are poorly developed or underused, inequitable distribution of health care workers within the governorates of the Gaza Strip and the focus is on the main governorate (Gaza) [14, 27, 28]. Nursing managers can implement systematic strategies to enhance nurses' knowledge and practice in BLS to target low-scoring Governorates.

Implication for decision makers and management

The study findings can enhance insight and understanding of the effects of the BLS training program in Nurses' knowledge. We suggest some practical implications for managers. Advanced BLS training for all healthcare providers, doctors, and nurses working in hospitals and healthcare centers. Training in BLS must be systematic and continuous to increase confidence in the use of CPR and potentially save lives. Then, it is needed to evaluate the impact and of these interventions. Future research should be expanded to include government hospitals in the West Bank, private institutions, non-governmental organizations and UNRWA.

Strengths and limitations

The current study is one of the few studies that evaluates the effect of BLS training on nurses' knowledge among nurses in government hospitals in the Gaza Strip, Palestine. The study provides important evidence of the positive effects of the BLS training program in improving nurses' knowledge. This study has several limitations including the use of a quasi-experimental design without a control group. The duration of the program was short, and it needed to enrich the educational program and improve the simulation laboratories in the Gaza Strip. In addition, no follow-up was conducted to measure the impact of the program on patient safety. Moreover, there are other limitations, which are that the study included

hospitals in the Gaza Strip only and did not include hospitals in the West Bank. It was conducted in government hospitals and did not include private institutions, non-governmental organizations and UNRWA.

Conclusion

This study provides significant evidence of the positive effects of the BLS training program in improving nurses' knowledge; we recommend advanced BLS training for all healthcare providers, doctors, and nurses working in hospitals and healthcare centers. Training in BLS must be systematic and continuous to increase confidence in the use of CPR and potentially save lives. Healthcare professionals must work together to ensure that these courses are successfully received and completed. Future research should target healthcare providers in emergency departments who are the first possible contact with patients who require CPR. In addition, nursing managers can implement systematic strategies to enhance nurses' knowledge in BLS to target low-scoring Governorates.

List of abbreviations

BLS	Basic Life Support
AHA	American Heart Association
CPR	Cardio Pulmonary Resuscitation
AED	Automated External Defibrillator

Acknowledgements

The authors want to thank the nurses who participated in the study. We also would like to thank the Clinical Research Development Unit of Tabriz Valiasr Hospital, Tabriz University of Medical Sciences, Tabriz, Iran for their assistance in this research.

Authors' contributions

Y.F, A.H.A, O.J.E, and E.K designed and conducted the study, performed the analysis and drafted the manuscript. Y.A.S, A.K, and A.A advised on the study design, facilitated data collection and revised the manuscript. Y.A.S, A.K, and Y.F helped in data collection. O.J.E, A.H.A, and E.K interpretation of data and revised the manuscript. All authors reviewed and approved the manuscript.

Funding

The authors have not received any funding.

Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

Permission was obtained from the Health Research Department of the Ministry of Health. The researchers initially gave the participants a full explanation of the research and its aims, and the participants signed the written informed consent form. All experimental protocols were approved by the ethical committee of the Ministry of Health (No. 1114555). Additionally, the study was conducted in accordance with the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

- ¹General Directorate of Nursing, Ministry of Health, Gaza, Palestine
- ²General Directorate of Mental Health, Ministry of Health, Gaza, Palestine
- ³Medical Services Directorate, Gaza Strip, Palestine
- ⁴College of Health Professions, Israa University, Gaza, Palestine
- ⁵Health Research Unit, Ministry of Health, Gaza, Palestine
- ⁶Faculty of Nursing, Midwifery Department, Islamic University of Gaza, Gaza, Palestine
- ⁷Clinical Research Development Unit of Tabriz Valiasr Hospital, Tabriz University of Medical Sciences, Tabriz, Iran

Received: 9 July 2023 / Accepted: 4 October 2023

Published online: 20 October 2023

References

1. Diwakar M, Tripathi A, Joshi K, Memoria M, Singh P. Latest trends on heart disease prediction using machine learning and image fusion. *Materials Today: Proceedings*. 2021;37:3213-8.
2. Beauchamp M, Tom O, Walters S, Khan I, Anna F. Trends in cardiopulmonary resuscitation: a four year review. *West Afr J Med*. 2013;19:25-30.
3. Duff JP, Topjian AA, Berg MD, Chan M, Haskell SE, Joyner BL et al. 2019 American Heart Association focused update on pediatric basic life support: an update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Pediatrics*. 2020;145(1).
4. Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation*. 2015;131(4):e29–e322.
5. Gabriel IO, Aluko JO. Theoretical knowledge and psychomotor skill acquisition of basic life support training programme among secondary school students. *World J Emerg Med*. 2019;10(2):81.
6. Greif R, Lockey AS, Conaghan P, Lippert A, De Vries W, Monsieurs KG, et al. European resuscitation council guidelines for resuscitation 2015: Sect. 10. Education and implementation of resuscitation. *Resuscitation*. 2015;95:288–301.
7. Berger C, Brinkrolf P, Ertmer C, Becker J, Friederichs H, Wenk M, et al. Combination of problem-based learning with high-fidelity simulation in CPR training improves short and long-term CPR skills: a randomised single blinded trial. *BMC Med Educ*. 2019;19:1-10.
8. Ihunanya OM, Oke Michael R, Babcock B, Amere LT. Knowledge, attitude and practice of cardiopulmonary resuscitation among nurses in Babcock University Teaching Hospital in Ilishan-Remo, Ogun State, Nigeria. *Int J Caring Sci*. 2020;13(3):1773-82.
9. Meakim C, Boese T, Decker S, Franklin AE, Gloe D, Lioce L, et al. Standards of best practice: Simulation standard I: terminology. *Clin Simul Nurs*. 2013;9(6):3-511.
10. Fluharty L, Hayes AS, Milgrom L, Malarney K, Smith D, Reklau MA, et al. A multisite, multi-academic track evaluation of end-of-life simulation for nursing education. *Clin Simul Nurs*. 2012;8(4):e135-e43.
11. Sok SR, Kim JA, Lee Y, Cho Y. Effects of a simulation-based CPR training program on knowledge, performance, and stress in clinical nurses. *J Continuing Educ Nurs*. 2020;51(5):225-32.
12. Lendahls L, Oscarsson MG. Midwifery students' experiences of simulation-and skills training. *Nurse Educ Today*. 2017;50:12-6.
13. Palestinian Central Bureau of Statistics. Preliminary Results of the Population, Housing and Establishments Census, 2017. February, 2018.
14. Albelbeisi AH, Albelbeisi A, El Bilbeisi AH, Taleb M, Takian A, Akbari-Sari A. Barriers toward the practice of healthy behaviors among patients with non-communicable diseases in Gaza Strip, Palestine. *SAGE Open Medicine*. 2021;9:20503121211029179.
15. Krejcie RV, Morgan DW. Determining sample size for research activities. *Educ Psychol Meas*. 1970;30(3):607-10.
16. Association AH. American Heart Association Guidelines for CPR and ECC [Internet]. Dallas: AHA; 2020.
17. NHCPS SaLb. BLS Practice Test 2023 [Available from: <https://nhcps.com/bls-practice-quiz/>].
18. Frost J. Weighted Average. Formula & Calculation Examples 2023 [Available from: <https://statisticsbyjim.com/basics/weighted-average>].

19. Wazonis AR. Simulation debriefing practices in traditional baccalaureate nursing programs: National survey results. *Clin Simul Nurs*. 2015;11(2):110–9.
20. Demirtas A, Guvenc G, Aslan Ö, Unver V, Basak T, Kaya C. Effectiveness of simulation-based cardiopulmonary resuscitation training programs on fourth-year nursing students. *Australasian Emerg Care*. 2021;24(1):4–10.
21. Onan A, Simsek N, Elcin M, Turan S, Erbil B, Deniz KZ. A review of simulation-enhanced, team-based cardiopulmonary resuscitation training for undergraduate students. *Nurse Educ Pract*. 2017;27:134–43.
22. Partiprajak S, Thongpo P. Retention of basic life support knowledge, self-efficacy and chest compression performance in Thai undergraduate nursing students. *Nurse Educ Pract*. 2016;16(1):235–41.
23. Sankar J, Vijayakanthi N, Sankar MJ, Dubey N. Knowledge and skill retention of in-service versus preservice nursing professionals following an informal training program in pediatric cardiopulmonary resuscitation: a repeated-measures quasiexperimental study. *BioMed research international*. 2013;2013.
24. Maurya A. Effectiveness of Simulation Teaching on neonatal resuscitation Skill Procedure among nursing students. *Int J Sci Res (IJSR)*. 2015;4(1).
25. Habibli T, Ghezeljeh TN, Haghani S. The effect of simulation-based education on nursing students' knowledge and performance of adult basic cardiopulmonary resuscitation: a randomized clinical trial. *Nurs Pract Today*. 2020.
26. Alnutaifi N. Knowledge and skills of cardiopulmonary resuscitation among critical care nurses in Kuwaiti Hospitals. *Am J Nurs Res*. 2021;9(2):64–70.
27. Albelbeisi AH, Albelbeisi A, El Bilbeisi AH, Takian A, Akbari-Sari A. Capacity of Palestinian primary health care system to prevent and control of non-communicable diseases in Gaza Strip, Palestine: A capacity assessment analysis based on adapted WHO-PEN tool. *Int J Health Plann Manag*. 2020;35(6):1412–25.
28. Giacaman R, Khatib R, Shabaneh L, Ramlawi A, Sabri B, Sabatinelli G, et al. Health status and health services in the occupied Palestinian territory. *The Lancet*. 2009;373(9666):837–49.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com